



COMPOSITION:

Each 1000 ml Contains

Dextrose	(USP) 25000	mg
Vitamin C	(USP) 50000	mg
Glycerin	(USP) 20000	mg
Potassium Chloride	(USP) 23500	mg
Magnesium Chloride	(USP) 19000	mg
Vitamin E	(USP) 1000	mg
Betain	(USP) 1000	mg





is recommended for all types of poultry to prevent dehydration, help birds in facing different types of stress and support production.



INGREDIENT	FEATURE	BENEFIT	USAGE
Potassium Chloride and Magnesium Chloride	<ul style="list-style-type: none"> Re-establish the ionic and hydroelectric balance in the digestive tract and blood 	<ul style="list-style-type: none"> Stabilise the blood pH Restore the functions of vital organs such as kidneys, heart, lungs and liver 	<ul style="list-style-type: none"> During different types of stress like handling, vaccination, traveling, starting phase, disease, medication, feed adaptation, debeaking and marketing
Betaine	<ul style="list-style-type: none"> Regulate osmotic pressure 	<ul style="list-style-type: none"> Ease the absorption of water and electrolytes 	<ul style="list-style-type: none"> Prevents dehydration in starting chicks, during heatstroke and diseases (like Enteritis)
Dextrose	<ul style="list-style-type: none"> Rapidly available energy 	<ul style="list-style-type: none"> Prevent hypoglycemia 	<ul style="list-style-type: none"> Giving energy to boost the birds performance
Vit C + Vit E	<ul style="list-style-type: none"> Natural antioxidant Aromatic and appetite stimulating substance Effective against thermal stress 	<ul style="list-style-type: none"> Protection against free radicals Compensate lack of appetite Mobilise or produce glucose required to maintain homeostasis 	<ul style="list-style-type: none"> Reduce the damage caused by oxidation Support production by improving feed intake Improve heat stress induced problems like poor immunity, reduce feed intake, increase body temperature, decrease weight gain, loss in carcass weight and mortality in birds



علامات:

♦ موسمیاتی دباؤ (شدید گرمی) ♦ پرندوں کا ہانپنا ♦ ڈی ہائیڈریشن ♦ جسم میں نمکیات کی کمی

خصوصیات:

♦ فوری توانائی بحال ♦ گرمی کی شدت ختم ♦ اعضائے ریہہ کے کام اور عمل کو برقرار رکھے۔

♦ برڈ کے جسم میں پانی کی مقدار کو فوراً پورا اور برقرار رکھے۔

♦ پولٹری برڈ اور باہر کے شدید درجہ حرارت میں فوری مطابقت۔

طریقہ استعمال:

FROST مرغیوں میں مختلف قسم کے دباؤ میں 1 ملی لیٹر فی 4 لیٹر پانی میں ملا کر 3 سے 4 دن تک استعمال کریں۔

FROST چوزوں کی آمد پر اور شدید گرمی میں 2 ملی لیٹر فی لیٹر پانی میں ملا کر 3 دن تک استعمال کریں۔

FROST اینٹی بائیوٹک کے استعمال کے بعد 3 دن تک استعمال کریں۔

FROST فلاک کو فروخت کرنے سے 2 دن قبل استعمال کریں۔

Packing:

5 Liter

Dosage:

1 ml Per 4 Liters
of Drinking Water



Marketed by:

merlin

Life Care

Shah Rukn-e-Alam
Multan- Pakistan



Manufactured By.

BOP Nutraceuticals
Plot # A2, Phase-1A, M3 Sahiwal Interchange,
Faisalabad Industrial Estate, Development &
Management Company (FIEDMC) Faisalabad.